



Adams G, Bamman MM. Characterization and regulation of mechanical loading-induced compensatory muscle hypertrophy. *Comprehensive Physiology*. 2012;2829:2970. [[PubMed](#)]

Atherton PJ, Etheridge T, Watt PW, Wilkinson D, Selby A, Rankin D, Smith K, Rennie MJ. Muscle full effect after oral protein: time-dependent concordance and discordance between human muscle protein synthesis and mTORC1 signaling. *Am J Clin Nutr*. 2010;92(5):1080–8. doi: 10.3945/ajcn.2010.29819. [[PubMed](#)] [[Cross Ref](#)]

Berardi JM, Price TB, Noreen EE, Lemon PW. Postexercise muscle glycogen recovery enhanced with a carbohydrate-protein supplement. *Med Sci Sports Exerc*. 2006;38(6):1106–13. doi: 10.1249/01.mss.0000222826.49358.f3. [[PubMed](#)] [[Cross Ref](#)]

Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol*. 1997;273(1 Pt 1):E122–9. [[PubMed](#)]

Biolo G, Williams BD, Fleming RY, Wolfe RR. Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes*. 1999;48(5):949–57. doi: 10.2337/diabetes.48.5.949. [[PubMed](#)] [[Cross Ref](#)]

Blomstrand E, Saltin B, Blomstrand E, Saltin B. Effect of muscle glycogen on glucose, lactate and amino acid metabolism during exercise and recovery in human subjects. *J Physiol*. 1999;514(1):293–302. doi: 10.1111/j.1469-7793.1999.293af.x. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Bodine SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R, Zlotchenko E, Scrimgeour A, Lawrence JC, Glass DJ, Yancopoulos GD. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol*. 2001;3(11):1014–9. doi: 10.1038/ncb1101-1014. [[PubMed](#)] [[Cross Ref](#)]

Bohe J, Low JF, Wolfe RR, Rennie MJ. Latency and duration of stimulation of human muscle protein synthesis during continuous infusion of amino acids. *J Physiol*. 2001;532(Pt 2):575–9. [[PMC free article](#)] [[PubMed](#)]

Borsheim E, Cree MG, Tipton KD, Elliott TA, Aarsland A, Wolfe RR. Effect of carbohydrate intake on net muscle protein synthesis during recovery from resistance exercise. *J Appl Physiol*. 2004;96(2):674–8. doi: 10.1152/jappphysiol.00333.2003. [[PubMed](#)] [[Cross Ref](#)]

Breen L, Phillips SM. Interactions between exercise and nutrition to prevent muscle waste during aging. *Br J Clin Pharmacol*. 2012. [Epub ahead of print] [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

- Burd NA, Tang JE, Moore DR, Phillips SM. Exercise training and protein metabolism: influences of contraction, protein intake, and sex-based differences. *J Appl Physiol*. 2009;106(5):1692–701. doi: 10.1152/jappphysiol.91351.2008. [[PubMed](#)] [[Cross Ref](#)]
- Camera DM, West DW, Burd NA, Phillips SM, Garnham AP, Hawley JA, Coffey VG. Low muscle glycogen concentration does not suppress the anabolic response to resistance exercise. *J Appl Physiol*. 2012;113(2):206–14. doi: 10.1152/jappphysiol.00395.2012. [[PubMed](#)] [[Cross Ref](#)]
- Candow DG, Chilibeck PD. Timing of creatine or protein supplementation and resistance training in the elderly. *Appl Physiol Nutr Metab*. 2008;33(1):184–90. doi: 10.1139/H07-139. [[PubMed](#)] [[Cross Ref](#)]
- Capaldo B, Gastaldelli A, Antonello S, Auletta M, Pardo F, Ciociaro D, Guida R, Ferrannini E, Sacca L. Splanchnic and leg substrate exchange after ingestion of a natural mixed meal in humans. *Diabetes*. 1999;48(5):958–66. doi: 10.2337/diabetes.48.5.958. [[PubMed](#)] [[Cross Ref](#)]
- Churchley EG, Coffey VG, Pedersen DJ, Shield A, Carey KA, Cameron-Smith D, Hawley JA. Influence of preexercise muscle glycogen content on transcriptional activity of metabolic and myogenic genes in well-trained humans. *J Appl Physiol*. 2007;102(4):1604–11. [[PubMed](#)]
- Coffey VG, Shield A, Canny BJ, Carey KA, Cameron-Smith D, Hawley JA. Interaction of contractile activity and training history on mRNA abundance in skeletal muscle from trained athletes. *Am J Physiol Endocrinol Metab*. 2006;290(5):E849–55. doi: 10.1152/ajpendo.00299.2005. [[PubMed](#)] [[Cross Ref](#)]
- Creer A, Gallagher P, Slivka D, Jemiolo B, Fink W, Trappe S. Influence of muscle glycogen availability on ERK1/2 and Akt signaling after resistance exercise in human skeletal muscle. *J Appl Physiol*. 2005;99(3):950–6. doi: 10.1152/jappphysiol.00110.2005. [[PubMed](#)] [[Cross Ref](#)]
- Cribb PJ, Hayes A. Effects of supplement timing and resistance exercise on skeletal muscle hypertrophy. *Med Sci Sports Exerc*. 2006;38(11):1918–25. doi: 10.1249/01.mss.0000233790.08788.3e. [[PubMed](#)] [[Cross Ref](#)]
- Denne SC, Liechty EA, Liu YM, Brechtel G, Baron AD. Proteolysis in skeletal muscle and whole body in response to euglycemic hyperinsulinemia in normal adults. *Am J Physiol*. 1991;261(6 Pt 1):E809–14. [[PubMed](#)]
- Dennis PB, Jaeschke A, Saitoh M, Fowler B, Kozma SC, Thomas G. Mammalian TOR: a homeostatic ATP sensor. *Science*. 2001;294(5544):1102–5. doi: 10.1126/science.1063518. [[PubMed](#)] [[Cross Ref](#)]
- Derave W, Lund S, Holman GD, Wojtaszewski J, Pedersen O, Richter EA. Contraction-stimulated muscle glucose transport and GLUT-4 surface content are dependent on glycogen content. *Am J Physiol*. 1999;277(6 Pt 1):E1103–10. [[PubMed](#)]
- Erschine RM, Fletcher G, Hanson B, Folland JP. Whey protein does not enhance the adaptations to elbow flexor resistance training. *Med Sci Sports Exerc*. 2012;44(9):1791–800. doi: 10.1249/MSS.0b013e318256c48d. [[PubMed](#)] [[Cross Ref](#)]
- Esmarck B, Andersen JL, Olsen S, Richter EA, Mizuno M, Kjaer M. Timing of postexercise protein intake is important for muscle hypertrophy with resistance training in elderly humans. *J Physiol*. 2001;535(Pt 1):301–11. [[PMC free article](#)] [[PubMed](#)]
- Fluckey JD, Vary TC, Jefferson LS, Farrell PA. Augmented insulin action on rates of protein synthesis after resistance exercise in rats. *Am J Physiol*. 1996;270(2 Pt 1):E313–9. [[PubMed](#)]
- Fox AK, Kaufman AE, Horowitz JF. Adding fat calories to meals after exercise does not alter glucose tolerance. *J Appl Physiol*. 2004;97(1):11–6. doi: 10.1152/jappphysiol.01398.2003. [[PubMed](#)] [[Cross Ref](#)]

Fujita S, Dreyer HC, Drummond MJ, Glynn EL, Volpi E, Rasmussen BB. Essential amino acid and carbohydrate ingestion before resistance exercise does not enhance postexercise muscle protein synthesis. *J Appl Physiol*. 2009;106(5):1730–9. doi: 10.1152/jappphysiol.90395.2008. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Gelfand RA, Barrett EJ. Effect of physiologic hyperinsulinemia on skeletal muscle protein synthesis and breakdown in man. *J Clin Invest*. 1987;80(1):1–6. doi: 10.1172/JCI113033. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Glynn EL, Fry CS, Drummond MJ, Dreyer HC, Dhanani S, Volpi E, Rasmussen BB. Muscle protein breakdown has a minor role in the protein anabolic response to essential amino acid and carbohydrate intake following resistance exercise. *Am J Physiol Regul Integr Comp Physiol*. 2010;299(2):R533–40. doi: 10.1152/ajpregu.00077.2010. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Goodman CA, Mayhew DL, Hornberger TA. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell Signal*. 2011;23(12):1896–906. doi: 10.1016/j.cellsig.2011.07.013. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Greenhaff PL, Karagounis LG, Peirce N, Simpson EJ, Hazell M, Layfield R, Wackerhage H, Smith K, Atherton P, Selby A, Rennie MJ. Disassociation between the effects of amino acids and insulin on signaling, ubiquitin ligases, and protein turnover in human muscle. *Am J Physiol Endocrinol Metab*. 2008;295(3):E595–604. doi: 10.1152/ajpendo.90411.2008. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Heslin MJ, Newman E, Wolf RF, Pisters PW, Brennan MF. Effect of hyperinsulinemia on whole body and skeletal muscle leucine carbon kinetics in humans. *Am J Physiol*. 1992;262(6 Pt 1):E911–8. [[PubMed](#)]

Hoffman JR, Ratamess NA, Tranchina CP, Rashti SL, Kang J, Faigenbaum AD. Effect of protein-supplement timing on strength, power, and body-composition changes in resistance-trained men. *Int J Sport Nutr Exerc Metab*. 2009;19(2):172–85. [[PubMed](#)]

Hulmi JJ, Kovanen V, Selanne H, Kraemer WJ, Hakkinen K, Mero AA. Acute and long-term effects of resistance exercise with or without protein ingestion on muscle hypertrophy and gene expression. *Amino Acids*. 2009;37(2):297–308. doi: 10.1007/s00726-008-0150-6. [[PubMed](#)] [[Cross Ref](#)]

Hulmi JJ, Lockwood CM, Stout JR. Effect of protein/essential amino acids and resistance training on skeletal muscle hypertrophy: A case for whey protein. *Nutr Metab (Lond)* 2010;7:51. doi: 10.1186/1743-7075-7-51. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Ivy J, Portman R. *Nutrient Timing: The Future of Sports Nutrition*. North Bergen, NJ: Basic Health Publications; 2004.

Ivy JL, Goforth HW Jr, Damon BM, McCauley TR, Parsons EC, Price TB. Early postexercise muscle glycogen recovery is enhanced with a carbohydrate-protein supplement. *J Appl Physiol*. 2002;93(4):1337–44. [[PubMed](#)]

Ivy JL. Glycogen resynthesis after exercise: effect of carbohydrate intake. *Int J Sports Med*. 1998;19(Suppl 2):S142–5. [[PubMed](#)]

Izumiya Y, Hopkins T, Morris C, Sato K, Zeng L, Viereck J, Hamilton JA, Ouchi N, LeBrasseur NK, Walsh K. Fast/Glycolytic muscle fiber growth reduces fat mass and improves metabolic parameters in obese mice. *Cell Metab*. 2008;7(2):159–72. doi: 10.1016/j.cmet.2007.11.003. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

- Jacinto E, Hall MN. Tor signalling in bugs, brain and brawn. *Nat Rev Mol Cell Biol.* 2003;4(2):117–26. doi: 10.1038/nrm1018. [[PubMed](#)] [[Cross Ref](#)]
- Jentjens R, Jeukendrup A. Determinants of post-exercise glycogen synthesis during short-term recovery. *Sports Med.* 2003;33(2):117–44. doi: 10.2165/00007256-200333020-00004. [[PubMed](#)] [[Cross Ref](#)]
- Jentjens RL, van Loon LJ, Mann CH, Wagenmakers AJ, Jeukendrup AE. Addition of protein and amino acids to carbohydrates does not enhance postexercise muscle glycogen synthesis. *J Appl Physiol.* 2001;91(2):839–46. [[PubMed](#)]
- Kawanaka K, Nolte LA, Han DH, Hansen PA, Holloszy JO. Mechanisms underlying impaired GLUT-4 translocation in glycogen-supercompensated muscles of exercised rats. *Am J Physiol Endocrinol Metab.* 2000;279(6):E1311–8. [[PubMed](#)]
- Kerksick C, Harvey T, Stout J, Campbell B, Wilborn C, Kreider R, Kalman D, Ziegenfuss T, Lopez H, Landis J, Ivy JL, Antonio J. International Society of Sports Nutrition position stand: nutrient timing. *J Int Soc Sports Nutr.* 2008;5:17. doi: 10.1186/1550-2783-5-17. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
- Kettelhut IC, Wing SS, Goldberg AL. Endocrine regulation of protein breakdown in skeletal muscle. *Diabetes Metab Rev.* 1988;4(8):751–72. doi: 10.1002/dmr.5610040805. [[PubMed](#)] [[Cross Ref](#)]
- Kim DH, Kim JY, Yu BP, Chung HY. The activation of NF-kappaB through Akt-induced FOXO1 phosphorylation during aging and its modulation by calorie restriction. *Biogerontology.* 2008;9(1):33–47. doi: 10.1007/s10522-007-9114-6. [[PubMed](#)] [[Cross Ref](#)]
- Koopman R, Beelen M, Stellingwerff T, Pennings B, Saris WH, Kies AK, Kuipers H, van Loon LJ. Coingestion of carbohydrate with protein does not further augment postexercise muscle protein synthesis. *Am J Physiol Endocrinol Metab.* 2007;293(3):E833–42. doi: 10.1152/ajpendo.00135.2007. [[PubMed](#)] [[Cross Ref](#)]
- Koopman R, Wagenmakers AJ, Manders RJ, Zorenc AH, Senden JM, Gorselink M, Keizer HA, van Loon LJ. Combined ingestion of protein and free leucine with carbohydrate increases postexercise muscle protein synthesis in vivo in male subjects. *Am J Physiol Endocrinol Metab.* 2005;288(4):E645–53. [[PubMed](#)]
- Kukuljan S, Nowson CA, Sanders K, Daly RM. Effects of resistance exercise and fortified milk on skeletal muscle mass, muscle size, and functional performance in middle-aged and older men: an 18-mo randomized controlled trial. *J Appl Physiol.* 2009;107(6):1864–73. doi: 10.1152/jappphysiol.00392.2009. [[PubMed](#)] [[Cross Ref](#)]
- Kumar V, Atherton P, Smith K, Rennie MJ. Human muscle protein synthesis and breakdown during and after exercise. *J Appl Physiol.* 2009;106(6):2026–39. doi: 10.1152/jappphysiol.91481.2008. [[PubMed](#)] [[Cross Ref](#)]
- Lambert CP, Flynn MG. Fatigue during high-intensity intermittent exercise: application to bodybuilding. *Sports Med.* 2002;32(8):511–22. doi: 10.2165/00007256-200232080-00003. [[PubMed](#)] [[Cross Ref](#)]
- Layman DK. Protein quantity and quality at levels above the RDA improves adult weight loss. *J Am Coll Nutr.* 2004;23(6 Suppl):631S–6S. [[PubMed](#)]
- Lemon PW, Mullin JP. Effect of initial muscle glycogen levels on protein catabolism during exercise. *J Appl Physiol.* 1980;48(4):624–9. [[PubMed](#)]

- Levenhagen DK, Gresham JD, Carlson MG, Maron DJ, Borel MJ, Flakoll PJ. Postexercise nutrient intake timing in humans is critical to recovery of leg glucose and protein homeostasis. *Am J Physiol Endocrinol Metab.* 2001;280(6):E982–93. [[PubMed](#)]
- Levine JA, Abboud L, Barry M, Reed JE, Sheedy PF, Jensen MD. Measuring leg muscle and fat mass in humans: comparison of CT and dual-energy X-ray absorptiometry. *J Appl Physiol.* 2000;88(2):452–6. [[PubMed](#)]
- MacDougall JD, Ray S, Sale DG, McCartney N, Lee P, Garner S. Muscle substrate utilization and lactate production. *Can J Appl Physiol.* 1999;24(3):209–15. doi: 10.1139/h99-017. [[PubMed](#)] [[Cross Ref](#)]
- McBride A, Ghilagaber S, Nikolaev A, Hardie DG. The glycogen-binding domain on the AMPK beta subunit allows the kinase to act as a glycogen sensor. *Cell Metab.* 2009;9(1):23–34. doi: 10.1016/j.cmet.2008.11.008. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
- Miller SL, Tipton KD, Chinkes DL, Wolf SE, Wolfe RR. Independent and combined effects of amino acids and glucose after resistance exercise. *Med Sci Sports Exerc.* 2003;35(3):449–55. doi: 10.1249/01.MSS.0000053910.63105.45. [[PubMed](#)] [[Cross Ref](#)]
- Moore DR, Robinson MJ, Fry JL, Tang JE, Glover EI, Wilkinson SB, Prior T, Tarnopolsky MA, Phillips SM. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr.* 2009;89(1):161–8. [[PubMed](#)]
- Norton LE, Layman DK, Bunpo P, Anthony TG, Brana DV, Garlick PJ. The leucine content of a complete meal directs peak activation but not duration of skeletal muscle protein synthesis and mammalian target of rapamycin signaling in rats. *J Nutr.* 2009;139(6):1103–9. doi: 10.3945/jn.108.103853. [[PubMed](#)] [[Cross Ref](#)]
- O’Gorman DJ, Del Aguila LF, Williamson DL, Krishnan RK, Kirwan JP. Insulin and exercise differentially regulate PI3-kinase and glycogen synthase in human skeletal muscle. *J Appl Physiol.* 2000;89(4):1412–9. [[PubMed](#)]
- Parkin JA, Carey MF, Martin IK, Stojanovska L, Febbraio MA. Muscle glycogen storage following prolonged exercise: effect of timing of ingestion of high glycemic index food. *Med Sci Sports Exerc.* 1997;29(2):220–4. doi: 10.1097/00005768-199702000-00009. [[PubMed](#)] [[Cross Ref](#)]
- Phillips SM, Van Loon LJ. Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.* 2011;29(Suppl 1):S29–38. [[PubMed](#)]
- Phillips SM. The science of muscle hypertrophy: making dietary protein count. *Proc Nutr Soc.* 2011;70(1):100–3. doi: 10.1017/S002966511000399X. [[PubMed](#)] [[Cross Ref](#)]
- Pitkanen HT, Nykanen T, Knuutinen J, Lahti K, Keinanen O, Alen M, Komi PV, Mero AA. Free amino acid pool and muscle protein balance after resistance exercise. *Med Sci Sports Exerc.* 2003;35(5):784–92. doi: 10.1249/01.MSS.0000064934.51751.F9. [[PubMed](#)] [[Cross Ref](#)]
- Power O, Hallihan A, Jakeman P. Human insulinotropic response to oral ingestion of native and hydrolysed whey protein. *Amino Acids.* 2009;37(2):333–9. doi: 10.1007/s00726-008-0156-0. [[PubMed](#)] [[Cross Ref](#)]
- Rasmussen BB, Tipton KD, Miller SL, Wolf SE, Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol.* 2000;88(2):386–92. [[PubMed](#)]
- Rennie MJ, Bohe J, Smith K, Wackerhage H, Greenhaff P. Branched-chain amino acids as fuels and anabolic signals in human muscle. *J Nutr.* 2006;136(1 Suppl):264S–8S. [[PubMed](#)]

- Richter EA, Derave W, Wojtaszewski JF. Glucose, exercise and insulin: emerging concepts. *J Physiol.* 2001;535(Pt 2):313–22. [[PMC free article](#)] [[PubMed](#)]
- Robergs RA, Pearson DR, Costill DL, Fink WJ, Pascoe DD, Benedict MA, Lambert CP, Zachweija JJ. Muscle glycogenolysis during differing intensities of weight-resistance exercise. *J Appl Physiol.* 1991;70(4):1700–6. [[PubMed](#)]
- Roy BD, Tarnopolsky MA. Influence of differing macronutrient intakes on muscle glycogen resynthesis after resistance exercise. *J Appl Physiol.* 1998;84(3):890–6. [[PubMed](#)]
- Staples AW, Burd NA, West DW, Currie KD, Atherton PJ, Moore DR, Rennie MJ, Macdonald MJ, Baker SK, Phillips SM. Carbohydrate does not augment exercise-induced protein accretion versus protein alone. *Med Sci Sports Exerc.* 2011;43(7):1154–61. doi: 10.1249/MSS.0b013e31820751cb. [[PubMed](#)] [[Cross Ref](#)]
- Tang JE, Manolagos JJ, Kujbida GW, Lysecki PJ, Moore DR, Phillips SM. Minimal whey protein with carbohydrate stimulates muscle protein synthesis following resistance exercise in trained young men. *Appl Physiol Nutr Metab.* 2007;32(6):1132–8. doi: 10.1139/H07-076. [[PubMed](#)] [[Cross Ref](#)]
- Tarnopolsky MA, Bosman M, Macdonald JR, Vandeputte D, Martin J, Roy BD. Postexercise protein-carbohydrate and carbohydrate supplements increase muscle glycogen in men and women. *J Appl Physiol.* 1997;83(6):1877–83. [[PubMed](#)]
- Timmons JA. Variability in training-induced skeletal muscle adaptation. *J Appl Physiol.* 2011;110(3):846–53. doi: 10.1152/jappphysiol.00934.2010. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
- Tipton KD, Elliott TA, Cree MG, Aarsland AA, Sanford AP, Wolfe RR. Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab.* 2007;292(1):E71–6. [[PubMed](#)]
- Tipton KD, Elliott TA, Cree MG, Wolf SE, Sanford AP, Wolfe RR. Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise. *Med Sci Sports Exerc.* 2004;36(12):2073–81. [[PubMed](#)]
- Tipton KD, Elliott TA, Ferrando AA, Aarsland AA, Wolfe RR. Stimulation of muscle anabolism by resistance exercise and ingestion of leucine plus protein. *Appl Physiol Nutr Metab.* 2009;34(2):151–61. doi: 10.1139/H09-006. [[PubMed](#)] [[Cross Ref](#)]
- Tipton KD, Ferrando AA, Phillips SM, Doyle D Jr, Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol.* 1999;276(4 Pt 1):E628–34. [[PubMed](#)]
- Tipton KD, Rasmussen BB, Miller SL, Wolf SE, Owens-Stovall SK, Petrini BE, Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab.* 2001;281(2):E197–206. [[PubMed](#)]
- Verdijk LB, Jonkers RA, Gleeson BG, Beelen M, Meijer K, Savelberg HH, Wodzig WK, Dendale P, van Loon LJ. Protein supplementation before and after exercise does not further augment skeletal muscle hypertrophy after resistance training in elderly men. *Am J Clin Nutr.* 2009;89(2):608–16. doi: 10.3945/ajcn.2008.26626. [[PubMed](#)] [[Cross Ref](#)]
- Willoughby DS, Stout JR, Wilborn CD. Effects of resistance training and protein plus amino acid supplementation on muscle anabolism, mass, and strength. *Amino Acids.* 2007;32(4):467–77. doi: 10.1007/s00726-006-0398-7. [[PubMed](#)] [[Cross Ref](#)]
- Wilson GJ, Layman DK, Moulton CJ, Norton LE, Anthony TG, Proud CG, Rupassara SI, Garlick PJ. Leucine or carbohydrate supplementation reduces AMPK and eEF2 phosphorylation and extends

postprandial muscle protein synthesis in rats. *Am J Physiol Endocrinol Metab.* 2011;301(6):E1236–42. doi: 10.1152/ajpendo.00242.2011. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]

Wojtaszewski JF, MacDonald C, Nielsen JN, Hellsten Y, Hardie DG, Kemp BE, Kiens B, Richter EA. Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising human skeletal muscle. *Am J Physiol Endocrinol Metab.* 2003;284(4):E813–22. [[PubMed](#)]

Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, Tarnopolsky MA, Phillips SM. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr.* 2012;108(10):1780–8. doi: 10.1017/S0007114511007422. [[PubMed](#)] [[Cross Ref](#)]

Zawadzki KM, Yaspelkis BB 3rd, Ivy JL. Carbohydrate-protein complex increases the rate of muscle glycogen storage after exercise. *J Appl Physiol.* 1992;72(5)